

Anterior T-Wave Inversion Does Not Convey Short-Term Sudden Death Risk

Kirchhof, Paulus; Fabritz, Larissa

DOI:

[10.1016/j.jacc.2016.11.011](https://doi.org/10.1016/j.jacc.2016.11.011)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Kirchhof, P & Fabritz, L 2017, 'Anterior T-Wave Inversion Does Not Convey Short-Term Sudden Death Risk: Inverted Is the New Normal', *Journal of the American College of Cardiology*, vol. 69, no. 1, pp. 10-12.
<https://doi.org/10.1016/j.jacc.2016.11.011>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Inverted is the new normal. Anterior T wave inversion does not convey short-term sudden death risk in young white adults.

Larissa Fabritz, MD ¹⁻³, Paulus Kirchhof, MD ¹⁻⁴

¹Institute of Cardiovascular Sciences, University of Birmingham, ²Department of Cardiology, University Hospital Birmingham Foundation NHS Trust, ³Department of Cardiology, Hospital of the University of Münster, Germany, ⁴Department of Cardiology, SWBH NHS Trust

Correspondence address:

Professor Paulus Kirchhof
Institute of Cardiovascular Sciences,
University of Birmingham,
Wolfson Drive
Birmingham, B15 2TT
United Kingdom
Telephone: ++44 121 414 7042
Fax: 44 121 4145788
e-mail: p.kirchhof@bham.ac.uk

Ventricular fibrillation leading to sudden arrhythmic death, one of the major causes of death in young adults (together with road accidents, suicide, and violence), is a devastating event for affected families. (1) The first symptomatic manifestation is often lethal, despite better dissemination of first responder support for witnessed sudden arrhythmic death events, automated defibrillators, and faster “chains of survival” supporting acute care. Hence, early detection of individuals at risk is needed to prevent sudden arrhythmic death in younger adults. Unfortunately, it is a bit like looking for a golden needle in the haystack (Figure 1).

Inherited cardiac conditions such as arrhythmogenic right ventricular cardiomyopathy (ARVC), hypertrophic cardiomyopathy, or long QT syndrome, are among the more common causes of sudden death in young adults. The ECG is often sufficient to detect long QT syndrome, and has a reasonable sensitivity to detect hypertrophic cardiomyopathy. ECG screening has been successfully applied to detect long QT syndrome and Brugada ECG patterns in population-based studies (2,3) including the very young. (4-6) Many soccer, basketball, baseball, and cricket clubs routinely require an ECG and other cardiac tests before they sign contracts with professional players. Several international organisations, including the International Olympic Committee and other expert groups have proposed ECG screening in young adults and athletes (7-9), but it is unclear whether all athletes should undergo ECG screening. (10,11)

Screening tests require a high specificity to minimize anxiety and resource-use for further work-up in carriers of a given marker. In this issue of JACC, Dr. Malhotra and colleagues report a systematic analysis of over 16000 ECGs of Caucasian young adults including almost 3000 athletes (12). They found that T wave inversions in leads V1 and V2 are a normal variant, present in 1.4% of men and 4.3% of women studied. T wave inversions beyond V2 (i.e. in V1-V3 or in more than three chest leads) was rare (0.2%) in males and uncommon (1.2%) in females (2.1% in female athletes). None of the probands studied fulfilled the diagnostic criteria ARVC. (13) Furthermore, none of the probands with T wave inversion in the anterior chest leads suffered from an adverse event during a follow-up time of almost two years. This can clearly help clinicians who are faced with young adults showing anterior T wave inversion. The data also provide robust information on the spectrum of anterior T wave morphologies in young Caucasian adults, and support an updated definition of the normal range in Europe, as already suggested by the “Seattle criteria” (14).

The ECG integrates electrical differences throughout the chest, reflecting the main electrical gradients in the heart. Hence, it is less sensitive to changes in the right ventricle than in the larger left ventricle. Because ARVC affects the right ventricle earlier and more severely than the left ventricle, it can be more challenging to detect signs of ARVC in the ECG, compared to predominantly left ventricular diseases. Based on the position of the heart in the chest and

the type of recording of the chest (Wilson) ECG leads, the anterior chest leads of the ECG (V1-V3) are most likely to reveal right ventricular abnormalities. Indeed, anterior T wave inversion is found in many ARVC patients. (14,15) The report by Dr. Malhotra and colleagues clearly underpins that anterior T wave inversion in the anterior chest leads was not associated with a high risk of adverse events within a few years (12). It is conceivable that ARVC develops over a longer follow-up period in some of the probands. The data also suggest that the post-pubertal female anatomy of the chest is more conducive to T wave inversion in the anterior chest leads than male anatomies.

Detection of arrhythmogenic right ventricular cardiomyopathy appears of special relevance in athletes as endurance training accelerates the progression of disease. (16-18) Should we thus combine abnormal T wave patterns with other ECG markers for ARVC, e.g. further T wave abnormalities (19), S waves (13,20), or late potentials in the signal averaged ECG (13,21,22)? Should we alternatively stop using the ECG to screen for ARVC in young adults? The report by Dr Malhotra provides robust information on the relevance of T wave inversion in the short term in young adults. It is sufficient to reassure affected individuals that there is no imminent danger. Further work is needed to develop better tools to screen for rare but potentially life threatening diseases like ARVC.

Funding&Acknowledgments

This work was supported by the European Union Grant Agreement No. 633196 [CATCH ME] to Drs. Kirchhof and Fabritz); British Heart Foundation (FS/13/43/30324 to Drs. Kirchhof and Fabritz); Leducq Foundation to Dr. Kirchhof.

Dr. Fabritz has received further institutional research grant support from DFG, MRC, and Gilead Inc. Dr. Kirchhof has received further research support from the German Centre for Heart Research and from several drug and device companies active in atrial fibrillation; and has received honoraria from several such companies. Drs. Fabritz and Kirchhof are listed as inventors on a patent (WO2015/140571) held by the University of Birmingham on genotype-specific antiarrhythmic drug therapy of atrial fibrillation.

References

1. Priori SG, Blomstrom-Lundqvist C, Mazzanti A et al. 2015 ESC Guidelines for the management of patients with ventricular arrhythmias and the prevention of sudden cardiac death: The Task Force for the Management of Patients with Ventricular Arrhythmias and the Prevention of Sudden Cardiac Death of the European Society of Cardiology (ESC). Endorsed by: Association for European Paediatric and Congenital Cardiology (AEPC). *Eur Heart J* 2015;36:2793-867.
2. Sinner MF, Pfeufer A, Perz S et al. Spontaneous Brugada electrocardiogram patterns are rare in the German general population: results from the KORA study. *Europace* 2009;11:1338-44.
3. Junttila MJ, Raatikainen MJ, Karjalainen J, Kauma H, Kesaniemi YA, Huikuri HV. Prevalence and prognosis of subjects with Brugada-type ECG pattern in a young and middle-aged Finnish population. *Eur Heart J* 2004;25:874-8.
4. Yoshinaga M, Ushinohama H, Sato S et al. Electrocardiographic screening of 1-month-old infants for identifying prolonged QT intervals. *Circ Arrhythm Electrophysiol* 2013;6:932-8.
5. Schwartz PJ, Montemerlo M, Facchini M et al. The QT interval throughout the first 6 months of life: a prospective study. *Circulation* 1982;66:496-501.
6. Schwartz PJ, Stramba-Badiale M, Segantini A et al. Prolongation of the QT interval and the sudden infant death syndrome. *N Engl J Med* 1998;338:1709-14.
7. Corrado D, Pelliccia A, Bjornstad HH et al. Cardiovascular pre-participation screening of young competitive athletes for prevention of sudden death: proposal for a common European protocol. Consensus Statement of the Study Group of Sport Cardiology of the Working Group of Cardiac Rehabilitation and Exercise Physiology and the Working Group of Myocardial and Pericardial Diseases of the European Society of Cardiology. *Eur Heart J* 2005;26:516-24.
8. Pelliccia A, Zipes DP, Maron BJ. Bethesda Conference #36 and the European Society of Cardiology Consensus Recommendations revisited a comparison of U.S. and European criteria for eligibility and disqualification of competitive athletes with cardiovascular abnormalities. *J Am Coll Cardiol* 2008;52:1990-6.
9. IOC. Sudden Cardiovascular Death in Sport. Lausanne Recommendations for Preparticipation Cardiovascular Screening. 2004.
10. Corrado D, Basso C, Pavei A, Michieli P, Schiavon M, Thiene G. Trends in sudden cardiovascular death in young competitive athletes after implementation of a preparticipation screening program. *Jama* 2006;296:1593-601.
11. Maron BJ, Haas TS, Murphy CJ, Ahluwalia A, Rutten-Ramos S. Incidence and causes of sudden death in U.S. college athletes. *J Am Coll Cardiol* 2014;63:1636-43.
12. Malhotra A, Dhutia H, Gati S et al. The prevalence and significance of anterior T wave inversion in white, young athletes and non-athletes. *JACC* 2016;in press.
13. Marcus FI, McKenna WJ, Sherrill D et al. Diagnosis of arrhythmogenic right ventricular cardiomyopathy/dysplasia: proposed modification of the Task Force Criteria. *Eur Heart J* 2010;31:806-14.
14. Drezner JA, Ackerman MJ, Anderson J et al. Electrocardiographic interpretation in athletes: the 'Seattle criteria'. *Br J Sports Med* 2013;47:122-4.
15. Schnell F, Riding N, O'Hanlon R et al. Recognition and significance of pathological T-wave inversions in athletes. *Circulation* 2015;131:165-73.
16. Fabritz L, Hoogendijk MG, Scicluna BP et al. Load-reducing therapy prevents development of arrhythmogenic right ventricular cardiomyopathy in plakoglobin-deficient mice. *J Am Coll Cardiol* 2011;57:740-50.
17. James CA, Bhonsale A, Tichnell C et al. Exercise Increases Age-Related Penetrance and Arrhythmic Risk in Arrhythmogenic Right Ventricular Dysplasia/Cardiomyopathy Associated Desmosomal Mutation Carriers. *J Am Coll Cardiol* 2013.

18. Sawant AC, Bhonsale A, te Riele AS et al. Exercise has a disproportionate role in the pathogenesis of arrhythmogenic right ventricular dysplasia/cardiomyopathy in patients without desmosomal mutations. *J Am Heart Assoc* 2014;3:e001471.
19. Calore C, Zorzi A, Sheikh N et al. Electrocardiographic anterior T-wave inversion in athletes of different ethnicities: differential diagnosis between athlete's heart and cardiomyopathy. *Eur Heart J* 2016;37:2515-27.
20. Nasir K, Bomma C, Tandri H et al. Electrocardiographic features of arrhythmogenic right ventricular dysplasia/cardiomyopathy according to disease severity: a need to broaden diagnostic criteria. *Circulation* 2004;110:1527-34.
21. Kamath GS, Zareba W, Delaney J et al. Value of the signal-averaged electrocardiogram in arrhythmogenic right ventricular cardiomyopathy/dysplasia. *Heart Rhythm* 2011;8:256-62.
22. Santangeli P, Dello Russo A, Pieroni M et al. Fragmented and delayed electrograms within fibrofatty scar predict arrhythmic events in arrhythmogenic right ventricular cardiomyopathy: results from a prospective risk stratification study. *Heart Rhythm* 2012;9:1200-6.